

LIFE CHANGES

WHAT YOU LOVE DOESN'T HAVE TO


NUBEQA[®]
(darolutamide) 300 mg
tablets

A guide to NUBEQA[®]

NUBEQA is pronounced new-BECCA



WHAT IS NUBEQA[®]?

NUBEQA[®] (darolutamide) is a prescription medicine used to treat adults with prostate cancer:

- that has not spread to other parts of the body and no longer responds to a medical or surgical treatment that lowers testosterone
- that has spread to other parts of the body and responds to a medical or surgical treatment that lowers testosterone
- that has spread to other parts of the body and responds to a medical or surgical treatment that lowers testosterone in combination with docetaxel.

It is not known if NUBEQA is safe and effective in females or children.

IMPORTANT SAFETY INFORMATION

Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you:

- have heart problems, high blood pressure, diabetes, or high amounts of fat or cholesterol in your blood (dyslipidemia)
- have a history of seizures, brain injury, stroke, or brain tumors
- have kidney or liver problems

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).

ABOUT NUBEQA®

NUBEQA is a prescription medicine that's approved to treat certain types of advanced prostate cancer



Tablets shown are not actual size

Interested in information about NUBEQA results or caregiver-specific resources?

Use the buttons below to directly navigate.

If your prostate cancer **has spread** to other parts of the body (metastatic) and **responds to ADT**,* learn about the NUBEQA clinical studies relevant to you.

[See results from these clinical studies](#)



If your prostate cancer **has not spread** to other parts of the body (non-metastatic) and **does not respond to ADT**, learn about the NUBEQA clinical study relevant to you.

[See results from this clinical study](#)



If you're a caregiver, head to [page 13](#) for support.

*ADT refers to a medical or surgical treatment that lowers testosterone.

IMPORTANT SAFETY INFORMATION (continued)

Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you (continued):

- are pregnant or plan to become pregnant. NUBEQA can cause harm to your unborn baby and loss of pregnancy (miscarriage)
- have a partner who may become pregnant. Males who have female partners who may become pregnant should use effective birth control (contraception) during treatment and for 1 week after the last dose of NUBEQA. Talk with your healthcare provider about birth control methods
- are breastfeeding or plan to breastfeed. It is not known if NUBEQA passes into breast milk

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets



HOW NUBEQA® WORKS

What makes prostate cancer cells grow?

Prostate cancer cells contain special proteins called androgen receptors. When the hormone testosterone attaches to these receptors, it can trigger processes that cause prostate cancer cells to grow.

How does NUBEQA delay prostate cancer growth?

NUBEQA is an androgen receptor blocker, which means **it attaches to the androgen receptors inside cancer cells and blocks testosterone from attaching**. By being designed to stop testosterone attaching, NUBEQA may help delay cancer cells from growing.



IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. NUBEQA may affect the way other medicines work and other medicines may affect how NUBEQA works.

You should not start or stop any medicine before you talk with the healthcare provider that prescribed NUBEQA.

Know the medicines you take. Keep a list of them with you to show to your healthcare provider and pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

TAKING NUBEQA®

NUBEQA is a pill

2 tablets, 2X a day with food

Follow your doctor's instructions while taking NUBEQA



How should I take NUBEQA?

- Take NUBEQA exactly as your doctor tells you
- Your doctor may change your dose, temporarily stop, or completely stop your treatment with NUBEQA if you have certain side effects. Do not stop taking your prescribed dose of NUBEQA without talking with your doctor first
- Swallow two tablets whole, 2 times a day, with food
- Unless otherwise advised by your doctor, continue on ADT* while taking NUBEQA
- If you take more NUBEQA than prescribed, call your doctor right away

What should I do if I miss a dose of NUBEQA?

- Take your prescribed dose as soon as you remember
- Do not take 2 doses together to make up for a missed dose

*ADT refers to a medical or surgical treatment that lowers testosterone.

IMPORTANT SAFETY INFORMATION (continued)

NUBEQA may cause serious side effects, including:

- **Heart disease.** Blockage of the arteries in the heart (ischemic heart disease) that can lead to death has happened in some people during treatment with NUBEQA. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with NUBEQA. Call your healthcare provider or get medical help right away if you get chest pain or discomfort at rest or with activity, or shortness of breath during your treatment with NUBEQA.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

NUBEQA® OFFERS OPTIONS FOR A PERSONALIZED TREATMENT DECISION

If you have **metastatic** prostate cancer that responds to ADT*



YOU AND YOUR DOCTOR MAY DECIDE BETWEEN 2 TREATMENT OPTIONS:

 **NUBEQA** + ADT

OR

 **NUBEQA** + ADT + CHEMO[†]

Dosing with chemotherapy for metastatic prostate cancer that responds to ADT

NUBEQA is not chemotherapy, but one of its approved uses is in combination with a type of chemotherapy called docetaxel for the treatment of metastatic prostate cancer that responds to ADT.

- Chemotherapy is given as an intravenous (IV) infusion in your arm once every 3 weeks for up to 6 cycles
- All 6 cycles of chemotherapy will be given within the first ~5 months of treatment with NUBEQA
- Your doctor may change, delay, or stop treatment with chemotherapy if needed
- Once chemotherapy is complete, treatment with NUBEQA can be continued as prescribed by your doctor

*ADT refers to a medical or surgical treatment that lowers testosterone.

†Chemotherapy=docetaxel.

IMPORTANT SAFETY INFORMATION (continued)

NUBEQA may cause serious side effects, including (continued):

- **Seizure.** Treatment with NUBEQA may increase your risk of having a seizure. You should avoid activities where a sudden loss of consciousness could cause serious harm to yourself or others. Tell your healthcare provider right away if you have loss of consciousness or seizure during treatment with NUBEQA.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).



NUBEQA®
(darolutamide) 300 mg tablets

PROVEN RESULTS WITH NUBEQA®

FOR MEN WITH METASTATIC PROSTATE CANCER THAT RESPONDS TO ADT*

NUBEQA was evaluated in two different clinical trials, with and without chemotherapy.†

This gives you and your doctor options to tailor your treatment plan to your needs.

Study 1

NUBEQA + ADT

NUBEQA

 **REDUCED THE RISK**
of cancer getting worse by
ALMOST HALF

In a clinical study, NUBEQA + ADT reduced the risk of progression‡ by 46% compared to ADT alone (median§ not reached vs 25 months).

At the time of analysis, 29% (128 out of 446) of men taking NUBEQA + ADT had cancer worsening via scans or were no longer living compared to 42% (94 out of 223) of men taking ADT alone.

Study 2

NUBEQA + ADT + CHEMO

NUBEQA helped men

LIVE LIFE LONGER

NUBEQA + ADT + chemotherapy extended overall survival by lowering the risk of death by 32% compared to ADT + chemotherapy (median§ not reached vs 48.9 months).

At the time of analysis, 35% (229 out of 651) of men taking NUBEQA + ADT + chemotherapy were no longer living compared to 46% (304 out of 654) of men taking ADT + chemotherapy.

See full results



*ADT refers to a medical or surgical treatment that lowers testosterone.

†Chemotherapy=docetaxel.

‡rPFS is measured by the cancer worsening via scans or the participant no longer living.

§Median is the middle value in a set of numbers, not the average.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of NUBEQA include:

- increase in liver function tests
- decreased white blood cells (neutropenia)
- feeling more tired than usual

The most common side effects of NUBEQA in combination with docetaxel include:

- constipation
- rash
- decreased appetite
- bleeding
- weight gain
- high blood pressure
- decreased red blood cells (anemia)
- high blood sugar levels
- decreased white blood cells
- increase in liver function tests
- low blood calcium levels

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

HOW NUBEQA® WAS STUDIED

IN MEN WITH METASTATIC PROSTATE CANCER THAT RESPONDS TO ADT*

Study 1

Clinical study **without** chemotherapy

NUBEQA + ADT



446 RECEIVED

NUBEQA



ADT



223 RECEIVED

ADT

A clinical study of 669 men evaluated radiographic progression-free survival in men taking NUBEQA + ADT compared to men taking ADT alone.

Study 2

Clinical study **with** chemotherapy†

NUBEQA + ADT + CHEMO



651 RECEIVED

NUBEQA



ADT



Chemotherapy

6 cycles every 3 weeks over <5 months



654 RECEIVED

ADT



Chemotherapy

6 cycles every 3 weeks over <5 months

A clinical study of 1305 men evaluated overall survival in men taking NUBEQA + ADT + chemotherapy compared to men taking ADT + chemotherapy alone.

*ADT refers to a medical or surgical treatment that lowers testosterone.

†Chemotherapy=docetaxel.

IMPORTANT SAFETY INFORMATION (continued)

NUBEQA may cause fertility problems in males, which may affect the ability to father children. Talk to your healthcare provider if you have concerns about fertility.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

PROVEN RESULTS WITH NUBEQA®

FOR MEN WITH **NON-METASTATIC PROSTATE CANCER THAT DOES NOT RESPOND TO ADT***

NUBEQA HELPED MEN LIVE **2X LONGER** WITHOUT CANCER SPREADING vs ADT alone

Men who were treated with NUBEQA + ADT lived more than twice as long without their cancer spreading to other parts of the body, compared with men who took ADT alone (median† 3.4 years vs 1.5 years).

At the time of analysis, 23% (221 out of 955) of men taking NUBEQA + ADT had cancer spreading or were no longer living compared to 39% (216 out of 554) of men taking ADT alone.

Clinical study in non-metastatic prostate cancer that does not respond to ADT



955 RECEIVED

NUBEQA



ADT



554 RECEIVED

ADT

A clinical study of 1509 men compared men taking NUBEQA + ADT to men taking ADT alone.

See full results



*ADT refers to a medical or surgical treatment that lowers testosterone.

†Median is the middle value in a set of numbers, not the average.

IMPORTANT SAFETY INFORMATION (continued)

These are not all the possible side effects of NUBEQA.

Call your healthcare provider for medical advice about side effects. You are encouraged to report side effects or quality complaints of products to the FDA by visiting www.fda.gov/medwatch or calling [1-800-FDA-1088](tel:1-800-FDA-1088).

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at NUBEQA.com/PI.



NUBEQA®
(darolutamide) 300 mg tablets

SIDE EFFECTS

IN MEN WITH METASTATIC PROSTATE CANCER THAT RESPONDS TO ADT*

NUBEQA® may cause serious side effects, including:

Heart disease

Blockage of the arteries in the heart (ischemic heart disease) that can lead to death has happened in some people during treatment with NUBEQA. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with NUBEQA. Call your healthcare provider or get medical help right away if you get chest pain or discomfort at rest or with activity, or shortness of breath during your treatment with NUBEQA.

Seizure

Treatment with NUBEQA may increase your risk of having a seizure. You should avoid activities where a sudden loss of consciousness could cause serious harm to yourself or others. Tell your healthcare provider right away if you have loss of consciousness or seizure during treatment with NUBEQA.

The most common side effects of NUBEQA include:



Increase in liver function tests



Decreased white blood cells (neutropenia)



Feeling more tired than usual

The most common side effects of NUBEQA in combination with docetaxel include:

- Constipation
- Rash
- Decreased appetite
- Bleeding
- Weight gain
- High blood pressure
- Decreased red blood cells (anemia)
- High blood sugar levels
- Decreased white blood cells
- Increase in liver function tests
- Low blood calcium levels

NUBEQA may cause fertility problems in males, which may affect their ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of NUBEQA.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

*ADT refers to a medical or surgical treatment that lowers testosterone.

IMPORTANT SAFETY INFORMATION

Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you:

- have heart problems, high blood pressure, diabetes, or high amounts of fat or cholesterol in your blood (dyslipidemia)

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at NUBEQA.com/PI.


NUBEQA®
(darolutamide) 300 mg tablets

SIDE EFFECTS

IN MEN WITH **NON-METASTATIC PROSTATE CANCER THAT DOES NOT RESPOND TO ADT***

NUBEQA® may cause serious side effects, including:

Heart disease

Blockage of the arteries in the heart (ischemic heart disease) that can lead to death has happened in some people during treatment with NUBEQA. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with NUBEQA. Call your healthcare provider or get medical help right away if you get chest pain or discomfort at rest or with activity, or shortness of breath during your treatment with NUBEQA.

Seizure

Treatment with NUBEQA may increase your risk of having a seizure. You should avoid activities where a sudden loss of consciousness could cause serious harm to yourself or others. Tell your healthcare provider right away if you have loss of consciousness or seizure during treatment with NUBEQA.

The most common side effects of NUBEQA include:



Increase in liver
function tests



Decreased white blood
cells (neutropenia)



Feeling more
tired than usual

NUBEQA may cause fertility problems in males, which may affect their ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of NUBEQA.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

*ADT refers to a medical or surgical treatment that lowers testosterone.

IMPORTANT SAFETY INFORMATION (continued)

Before taking NUBEQA, tell your healthcare provider about all your medical conditions, including if you:

- have a history of seizures, brain injury, stroke, or brain tumors
- have kidney or liver problems
- are pregnant or plan to become pregnant. NUBEQA can cause harm to your unborn baby and loss of pregnancy (miscarriage)
- have a partner who may become pregnant. Males who have female partners who may become pregnant should use effective birth control (contraception) during treatment and for 1 week after the last dose of NUBEQA. Talk with your healthcare provider about birth control methods
- are breastfeeding or plan to breastfeed. It is not known if NUBEQA passes into breast milk

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at NUBEQA.com/PI.


NUBEQA®
(darolutamide) 300 mg
tablets

STORIES FROM REAL PEOPLE

//

I think second opinions are just necessary—assuring yourself that the path you’re going to take in treatment is the right path to go on.”

- **Steve**, real person taking NUBEQA®



//

For me, having the right support system in place made the biggest difference.”

- **Ryan**, real person taking NUBEQA



**Hear more stories from
men taking NUBEQA
and their caregivers**

[Visit our website >](#)



Have a story of your own to share?

Sign up to share your NUBEQA story, or call **1-800-288-8374**
to share your story with a live Bayer representative.

Sign up >

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. NUBEQA may affect the way other medicines work and other medicines may affect how NUBEQA works.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

SUPPORT & RESOURCES

Support for you while taking NUBEQA®

Access Services by Bayer™ can help you with information about paying for NUBEQA. To speak with a specialist, call **1-800-288-8374**, available Monday - Friday, 8 AM - 8 PM (ET).

IF YOU ARE NEW TO NUBEQA

Ask your provider about the 1-month Free Trial Program.



IF YOU HAVE COMMERCIAL INSURANCE

You may be eligible for the NUBEQA
\$0 Co-Pay Program.



IF YOU HAVE MEDICARE OR MEDICAID

You may qualify for help paying out-of-pocket costs for NUBEQA.

Government-funded assistance may be available. Check your eligibility at [medicare.gov/your-medicare-costs](https://www.medicare.gov/your-medicare-costs).

Independent foundations may also be able to help.

CancerCare®
1-800-813-4673

HealthWell Foundation®
1-800-675-8416

PAN Foundation
1-866-316-7263

The Assistance Fund
1-855-845-3663

Patient Advocate Foundation Co-Pay Relief
1-866-512-3861

IF YOU DO NOT HAVE INSURANCE

Contact the Bayer US **Patient Assistance Foundation**, a charitable organization that helps eligible patients get Bayer prescription medicine at no cost.

Call **1-866-228-7723** Monday - Friday 9:00 AM - 6:00 PM (ET) or visit the foundation website at patientassistance.bayer.us to see if you qualify.

Depending on insurance plans and other factors, the cost of NUBEQA may differ from person to person. To learn about your insurance coverage and paying for NUBEQA, call **1-800-288-8374** to speak with a specialist at Access Services by Bayer.

IMPORTANT SAFETY INFORMATION (continued)

You should not start or stop any medicine before you talk with the healthcare provider that prescribed NUBEQA.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).


NUBEQA®
(darolutamide) 300 mg tablets

CAREGIVERS NEED CARE, TOO

Resources made specifically for caregivers

As a caregiver, it's important to take care of yourself as well as others. The following resources offer tips and support designed specifically for you.



My advice to caregivers would be to accept help from their family members, their loved ones, and even from the the person that they're taking care of."

- **Nancy**, former nurse and caregiver to a real person taking NUBEQA®



Tips for caregivers



Focus on your wellness with a healthy diet and exercise.



Ask for help from family and friends in caring for your loved one.



Do things you enjoy outside the home, such as meeting a friend for lunch or taking a walk.

Get more tips for caregivers on our [website](#) >

Hotlines

CancerCare®

Call **1-800-813-HOPE (4673)**

Monday - Thursday 10:00 AM - 6:00 PM (ET)
Friday 10:00 AM - 5:00 PM (ET)

Cancer Support Community

Call **1-888-793-9355**
for mental health support

AARP

Call **1-877-333-5885** (English),
or **1-888-971-2013** (Spanish)
Monday - Friday, 8:00 AM - 8:00 PM (ET)

Prostate cancer support groups

These organizations can help you find local support groups, online peer support, and tips for caregivers of men with prostate cancer.

ZERO Prostate Cancer

Prostate Cancer Foundation

Prostate Cancer Research Institute

FOR MEN TAKING NUBEQA: THANK YOUR CAREGIVER

Select and personalize a thank-you card for those who support you by downloading an option from [our website](#) >

IMPORTANT SAFETY INFORMATION (continued)

Know the medicines you take. Keep a list of them with you to show to your healthcare provider and pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout, Important Facts about NUBEQA on pages [14-15](#), and full Prescribing Information at [NUBEQA.com/PI](#).


NUBEQA®
(darolutamide) 300 mg tablets

IMPORTANT FACTS ABOUT NUBEQA® (darolutamide) tablets

Pronounced new-BECCA

WHAT IS NUBEQA?

NUBEQA is a prescription medicine used to treat adults with prostate cancer:

- that has not spread to other parts of the body and no longer responds to a medical or surgical treatment that lowers testosterone
- that has spread to other parts of the body and responds to a medical or surgical treatment that lowers testosterone
- that has spread to other parts of the body and responds to a medical or surgical treatment that lowers testosterone in combination with docetaxel.

It is not known if NUBEQA is safe and effective in females or children.

WHAT SHOULD I TELL MY HEALTHCARE PROVIDER BEFORE TAKING NUBEQA?

Tell your healthcare provider about all your medical conditions, including if you:

- have heart problems
- have high blood pressure
- have diabetes
- have high amounts of fat or cholesterol in your blood (dyslipidemia)
- have a history of seizures, brain injury, stroke, or brain tumors
- have kidney or liver problems
- are pregnant or plan to become pregnant. NUBEQA can cause harm to your unborn baby and loss of pregnancy (miscarriage).
- have a partner who may become pregnant. Males who have female partners who may become pregnant should use effective birth control (contraception) during treatment and for 1 week after the last dose of NUBEQA. Talk with your healthcare provider about birth control methods.
- are breastfeeding or plan to breastfeed. It is not known if NUBEQA passes into breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. NUBEQA may affect the way other medicines work and other medicines may affect how NUBEQA works.

You should not start or stop any medicine before you talk with the healthcare provider that prescribed NUBEQA.

Know the medicines you take. Keep a list of them with you to show to your healthcare provider and pharmacist when you get a new medicine.

HOW SHOULD I TAKE NUBEQA?

- Take NUBEQA exactly as your healthcare provider tells you.
- Your healthcare provider may change your dose, temporarily stop, or completely stop your treatment with NUBEQA if you have certain side effects.
- Take your prescribed dose of NUBEQA 2 times a day with food.
- Swallow NUBEQA tablets whole.
- If you are receiving gonadotropin-releasing hormone (GnRH) therapy, you should continue with this treatment during your treatment with NUBEQA unless you have had a surgery to lower the amount of testosterone in your body (surgical castration).
- If you miss a dose of NUBEQA, take your prescribed dose as soon as you remember before the next scheduled dose. Do not take 2 doses together to make up for a missed dose.
- If you take more NUBEQA than prescribed, call your healthcare provider right away.

Important Facts about NUBEQA are continued on the next page.

Please see full Prescribing Information at [NUBEQA.com/PI](https://www.nubeqa.com/PI).



IMPORTANT FACTS ABOUT NUBEQA® (darolutamide) tablets (cont'd)

Pronounced new-BECCA

WHAT ARE THE POSSIBLE SIDE EFFECTS OF NUBEQA?

NUBEQA may cause serious side effects, including:

- **Heart disease.** Blockage of the arteries in the heart (ischemic heart disease) that can lead to death has happened in some people during treatment with NUBEQA. Your healthcare provider will monitor you for signs and symptoms of heart problems during your treatment with NUBEQA. Call your healthcare provider or get medical help right away if you get chest pain or discomfort at rest or with activity, or shortness of breath during your treatment with NUBEQA.
- **Seizure.** Treatment with NUBEQA may increase your risk of having a seizure. You should avoid activities where a sudden loss of consciousness could cause serious harm to yourself or others. Tell your healthcare provider right away if you have loss of consciousness or seizure during treatment with NUBEQA.

The most common side effects of NUBEQA include:

- increase in liver function tests
- decreased white blood cells (neutropenia)
- feeling more tired than usual

The most common side effects of NUBEQA in combination with docetaxel include:

- | | |
|-----------------------|--------------------------------------|
| • constipation | • decreased red blood cells (anemia) |
| • rash | • high blood sugar levels |
| • decreased appetite | • decreased white blood cells |
| • bleeding | • increase in liver function tests |
| • weight gain | • low blood calcium levels |
| • high blood pressure | |

NUBEQA may cause fertility problems in males, which may affect the ability to father children.

Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of NUBEQA.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

HOW SHOULD I STORE NUBEQA?

- Store NUBEQA at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep the bottle tightly closed after you first open it.

Keep NUBEQA and all medicines out of the reach of children.

THE INFORMATION PROVIDED HERE IS NOT COMPREHENSIVE. TO LEARN MORE ABOUT NUBEQA:

- Talk to your healthcare provider or pharmacist
- Visit NUBEQA.com to obtain the FDA-approved product labeling
- Call Bayer at 1-888-842-2937

Please see full Prescribing Information at NUBEQA.com/PI.


NUBEQA®
(darolutamide) 300 mg
tablets

To learn more about NUBEQA[®],
talk to your doctor or visit [NUBEQA.com](https://www.nubeqa.com)



© 2025 Bayer. All rights reserved. BAYER, the Bayer Cross and NUBEQA are registered trademarks of Bayer, and Access Services by Bayer is a trademark of Bayer.
PP-NUB-US-4074-1 06/25


NUBEQA[®]
(darolutamide) 300 mg
tablets